

## COUNTY OF SAN DIEGO HEALTH & HUMAN SERVICES AGENCY EMERGENCY MEDICAL SERVICES



### COUNTY OF SAN DIEGO EMERGENCY MEDICAL SERVICES POLICY/PROCEDURE/PROTOCOL

SUBJECT: PEDIATRIC TREATMENT PROTOCOL ALTERED NEUROLOGIC FUNCTION (NON-TRAUMATIC)

No. <u>S-161</u> Page: <u>1 of 1</u>

Date: 07/01/2018

### BLS ALS

- Ensure patent airway, O<sub>2</sub> and/or ventilate prn
- O<sub>2</sub> Saturation
- Spinal stabilization when indicated
- Secretion problems; position on affected side
- Do not allow patient to walk
- Restrain prn
- Monitor blood glucose prn

# Hypoglycemia (suspected) or patient's glucometer results, if available, read <60 mg/dL (Neonate <45 mg/dL):

- If patient is awake and has gag reflex, give oral glucose paste or 3 tablets (15 g).
   Patient may eat or drink if able.
- If patient is unconscious, NPO.

#### Seizures:

- Protect airway, and protect from injury.
- Treat associated injuries.
- If febrile, remove excess clothing/covering.

### **Behavioral Emergencies:**

- Restrain only if necessary to prevent injury.
- Avoid unnecessary sirens.
- Consider law enforcement support.

- IV SO adjust prn
- Monitor EKG /blood glucose prn
- Capnography SO prn

### <u>Symptomatic ?opioid OD (excluding opioid dependent pain management patients)</u>:

o Naloxone per drug chart IN/IV/IM SO. MR SO

### Symptomatic ?opioids OD in opioid dependent pain management patients:

○ Naloxone titrate per drug chart IV (dilute IV dose per drug chart) or IN/IM per drug chart SQ. MR BH©

### Hypoglycemia:

Symptomatic patient unresponsive to oral glucose agents:

- o D₁₀ per drug chart IV <u>SO</u> if BS <60 mg/dL (Neonate <45 mg/dL)
- $_{\odot}$  If patient remains symptomatic and BS remains <60 mg/dL (Neonate <45 mg/dL) MR  $\underline{SO}$
- If no IV: Glucagon per drug chart IM <u>SO</u> if BS <60 mg/dL (Neonate <45 mg/dL)</li>

### Seizures:

### For:

- A. Ongoing generalized seizure lasting ≥5 minutes (includes seizure time prior to arrival of prehospital provider) SO
- B. Partial seizure with respiratory compromise SO
- C. Recurrent tonic-clonic seizures without lucid interval SO

#### GIVE:

 $\circ$  Versed per drug chart slow IV, (d/c if seizure stops)  $\underline{SO}.$  MR x1 in 10 minutes  $\underline{SO}$ 

#### If no IV:

o Versed per drug chart IN/IM SO. MR x1 in 10 minutes SO